

TYPE OF REPORT: Semiannual

TIME PERIOD: Jan-June, 1995

NAME AND LOCATION: Alan Strahler, Boston University

CONTRACT NUMBER: NAS5-31369

## ABSTRACT -- KEY POINTS

The semiempirical models for BRDF function fitting continue to acquit themselves well when tested on BRF measurements from various land cover types. The Li-Strahler mutual shadowing model, which provides the basis for two of the semiempirical kernels, fits both lab and field data from the CASI instrument well. Both albedo and semiempirical-parameter retrievals are not overly sensitive to noise in input measurements. The BRDF/Albedo product ATBD was completely rewritten and made available to the project on May 15. Coding for the Beta-3 delivery began late in the period.

Algorithm development for the land cover and land-cover change products continued with the assembly of a multirate TM database for the Walnut Gulch region of Arizona. Further data development awaits the receipt of better ground truth maps. Possible feature selection strategies were reviewed in the literature. Applications of neural net classifiers to the mixed pixel problem and to change detection were explored. Coding for the Beta-3 delivery began late in the period.

## TASK PROGRESS

### BRDF/Albedo Product

#### Model Development

During the reporting period, we continued our development of semiempirical models of bidirectional reflectance. This culminated with the submission of a paper to the Journal of Geophysical Research documenting the mathematical derivation and properties of the semiempirical kernels.

#### Model Validation

We also continued BRDF model validation activities by further work with the bidirectional reflectance data for eleven land cover types as provided by Dan Kimes of Goddard Space Flight Center. We also fit the semiempirical models to new suites of data acquired under our direction in July, 1994, at the Solar Simulation Laboratory, Changchun, China. The results were summarized in a proceedings paper for presentation at IGARSS'95 in July.

Additional validation efforts involved collaboration with John Miller and Ray Soffer, at York University, on a comparison of the Li-Strahler mutual shadowing model with observations using the CASI sensor made in Miller's laboratory on a forest of artificial trees. The model acquitted itself well. The results were written up for publication under joint authorship as a proceedings paper for presentation at IGARSS'95 in July.

In a collaboration with Robert d'Entremont of USAF Phillips Labs, we developed a plan for acquisition of cloud-cleared AVHRR data for the New England region with the objective of testing the ability of the Ambrals semiempirical models to capture the shape of land BRDFs over large regions. This will be largely top-of-atmosphere data, but atmospheric correction may also be attempted.

## Software and Algorithm Development

During the second quarter of the reporting period, the development of beta-delivery software began in earnest. This required major investments of science time in coding the algorithm and dealing with the PGS toolkit. During this time, Ambrals was also recoded and rereleased to be consistent with the MODIS algorithm.

## Atmospheric Correction-BRDF Effects

In collaboration with Eric Vermote, we continued developing the BRDF-atmospheric correction loop plan for MODIS, including the concept of Level 2G data -- forward binned, but unresampled values placed into the Level 3 grid.

## Sensitivity Studies

Using a theoretical approach, we explored the sensitivity of the retrieval of semiempirical model parameters and albedo to noise in the directional observations. We found that for a noise level of 5 percent, we found that parameters were retrieved with 1-4 percent accuracy for practical cases and albedo was retrieved with accuracies less than 2 percent.

## ATBD Revision

The ATBD document was totally rewritten during a four-month period, and the revised document (Version 3.2) was released to the project on May 15.

## Land Cover/Land-Cover Change

During this reporting period, we focused primarily on algorithm development for land cover classification. We continued our work with advanced technology (AT) classifiers -- neural nets, decision trees, adaptive classifiers, etc.

## Walnut Gulch

We received a series of TM images and some ground truth information on the Walnut Gulch site from Team Member Alfredo Huete's group, and spent considerable time registering and radiometrically correcting a time sequence of six selected TM datasets. We then began an exploration of neural network and tree-base classifiers. However, the quality of the ground truth information was limiting, and a new set of ground truth data was expected to be available shortly. Thus, we halted work with this dataset, pending the receipt of a new land cover map for the area.

## Feature Selection

A problem with MODIS data is that its multiple information domains provide a very large number of variables -- "features" -- that are observed for each pixel over the course of an annual cycle. This information must be reduced and stripped of redundancy for a classifier to use it effectively. An important activity in the latter part of the reporting period was a survey of the artificial intelligence and remote sensing literatures for fast and effective techniques for feature selection.

## MAS-BOREAS Data

We requested the navigated and resampled MODIS Airborne Simulator (MAS) data from BOREAS from BORIS, but were unable to secure it during the reporting period. The objective was

to provide a land-cover type classification using advanced-technology classifiers from MAS imagery and ground truth provided by the BORIS.

#### Neural Nets

Research on neural net classifiers continued with completion of a study on using neural net classifier outputs to indicate mixed pixels. In other work, fuzzy ARTMAP was used successfully for change detection in a database of TM imagery for the Tahoe National Forest.

#### Algorithm Coding

Coding of the land-cover algorithm required a major effort during the period. Initial coding strategies were developed late in the reporting interval.

### ANTICIPATED ACTIVITIES DURING THE NEXT QUARTER

#### BRDF/Albedo Product

Our primary activity during the next quarter for the BRDF/Albedo product will be to code the Beta-3 version of the algorithm, integrating the Toolkit and related subroutines. We will also continue refining the application of the semiempirical models, their validation, and their use to calculate albedo in realistic MODIS-MISR sensing scenarios.

#### Land Cover/Land-cover Change Product

During the next quarter, we will continue to code the Beta-3 version of the land-cover algorithm in the ESDIS environment. This will include both the monthly algorithm, which prepares a monthly database, and the classifier itself, which reads a year of monthly databases and provides a classified product every three months. In land-cover change activities, we will continue our liaison with Eric Lambin at the European Economic Community's Joint Research Center. Activities there focus on land-cover change as inferred from a ten year record of African AVHRR observations.

### PROBLEMS/CORRECTIVE ACTIONS

During this reporting period, we did not encounter any significant problems requiring corrective actions beyond the everyday problems that occur in research and algorithm development.

### PUBLICATIONS

Attached below is an updated list of publications derived with full or partial support under this contract. Items marked with an asterisk are new publications that are being submitted separately in hardcopy to the project management.

#### MASTER MODIS PUBLICATION LIST

July 1, 1995

Submitted

Schaaf, C. B. and A. Strahler, 1996, Modeling the hemispherical reflectances of a mountainous forest landscape using geometric-optics and a distributed parameter approach, submitted to Remote Sens. Env. (May, 1995)

## In Press

Strahler, A. H., 1995, Vegetation canopy reflectance modeling -- Recent developments and remote sensing perspectives, *Remote Sens. Rev.*, in press.

Wanner, W., X. Li, and A. H. Strahler, 1996, On the derivation of kernels for kernel-driven models of bidirectional reflectance, *J. Geophys. Res.*, in press.

Moody, A. and Woodcock, C. E., 1995, The influence of the spatial characteristics of landscapes on land-cover mapping using remote sensing, *Landscape Ecology*, in press.

Woodcock, C. E., J. B. Collins, and D. L. B. Jupp, 1995, Scaling remote sensing models, in *Scaling Up*, P. van Gardingen, G. Foody, and P. Curran (eds.), Society of Experimental Biology, Cambridge University Press, in press.

Wang, J., X. Li, and A. Strahler, 1995, Tree tomography--an indirect method for measuring crown structure, *Proc. of GeoInformatics'95*, in press.

Strahler, A., W. Wanner, Q. Zhu, and X. Jin, 1995, Bidirectional reflectance modeling of data from vegetation obtained in the Changchun solar simulation laboratory, *Proc. 15th Int. Geosci. and Remote Sensing Symp.*, Florence, Italy, July 10-14, 1995, in press.

Soffer, R. J. Miller, W. Wanner, and A. Strahler, 1995, Winter boreal forest canopy BRF results: Comparisons between airborne data, laboratory simulations and geometrical-optical model data, *Proc. 15th Int. Geosci. and Remote Sensing Symp.*, Florence, Italy, July 10-14, 1995, in press.

Strahler, A., A. Moody and E. Lambin, 1995, Land cover and land-cover change from MODIS, *Proc. 15th Int. Geosci. and Remote Sensing Symp.*, Florence, Italy, July 10-14, 1995, in press.

Wanner, W., A. Strahler, J.-P. Muller, M. Barnsley, P. Lewis, X. Li and C. Barker Schaaf, 1995, Global mapping of bidirectional reflectance and albedo for the EOS MODIS project: The algorithm and the product, *Proc. 15th Int. Geosci. and Remote Sensing Symp.*, Florence, Italy, July 10-14, 1995, in press.

Wanner, W., X. Li, and A. Strahler, 1995, A new class of geometric-optical semiempirical kernels for global BRDF and albedo modeling, *Proc. 15th Int. Geosci. and Remote Sensing Symp.*, Florence, Italy, July 10-14, 1995, in press.

Liang, S. and A. H. Strahler, 1995, An analytic radiative transfer model for a coupled atmosphere and leaf canopy, *J. Geophys. Res.*, in press.

Li, X., A. H. Strahler, and C. E. Woodcock, 1995, A hybrid geometric optical-radiative transfer approach for modeling albedo and directional reflectance of discontinuous canopies, *IEEE Trans. Geosci. and Remote Sens.*, in press.

## Published

\*Li, X., R. Davis, and C. Woodcock, 1994, A hybrid geometric optical and radiative transfer approach for modeling pyranometer measurements under a jack pine forest, *Geographic Information Sci.*, vol. 1, pp. 27-34.

\*Moody, A. and Woodcock, C. E., 1995, Calibration-based methods for correcting coarse resolution estimates of land-cover proportions, *Proc. 61st Ann. Convention Amer. Soc. Photogrammetry and Remote Sens.*, Charlotte, NC, Feb. 27-March 2, 1995, vol. 3, pp. 684-693.

\*Borak, J., P. Fisher, A. Strahler, and A. Moody, 1995, Local-scale evaluation of a technique for land-cover classification based on composited NDVI data, Proc. 61st Ann. Convention Amer. Soc. Photogrammetry and Remote Sens., Charlotte, NC, Feb. 27-March 2, 1995, vol. 3, pp. 796-805.

\*Li, X. A. H. Strahler, and C. E. Woodcock, 1994, Hybrid geometric-optical radiative-transfer model for the directional reflectance of discontinuous vegetation canopies, Proc. European Symp. on Satellite Remote Sensing, Sept. 26-30, Rome, Italy (Eur. Optical Soc., SPIE-Int. Soc. for Optical Engr.), pp. 27-37.

\*Schaaf, C. B. and A. H. Strahler, 1994, Using a geometric-optical model to calculate the bidirectional and hemispherical reflectance of forested slopes, Proc. European Symp. on Satellite Remote Sensing, Sept. 26-30, Rome, Italy (Eur. Optical Soc., SPIE-Int. Soc. for Optical Engr.), pp. 21-26.

\*Running, S., C. Justice, D. Hall, A. Huete, Y. Kaufmann, J-P. Muller, A. Strahler, V. Vanderbilt, Z-M. Wan, P. Teillet and D. Carnegie, 1994, Terrestrial remote sensing science and algorithms planned for EOS/MODIS, Int. J. Remote Sens., vol. 15, pp. 3587-3620.

\*Moody, A. and A. H. Strahler, 1994, Characteristics of composited AVHRR data and problems in their classification, Int. J. Remote Sens., vol. 15, pp. 3473-3491.

Schaaf, C. B., X. Li and A. H. Strahler, 1994, Topographic effects of bidirectional and hemispherical reflectances calculated with a geometric-optical canopy model, IEEE Trans. Geosci. and Remote Sens., vol. 32, pp. 1186-1193.

Schaaf, C. B. and A. H. Strahler, 1994, Validation of bidirectional and hemispherical reflectances from a geometric-optical model using ASAS Imagery and Pyranometer Measurements of a Spruce Forest, Remote Sens. Environ., vol. 49, pp. 138-144.

Gopal, S, D. M. Sklarew and E. Lambin, 1994, Fuzzy-neural networks in multitemporal classification of landcover change in the Sahel, in New Tools for Spatial Analysis: Proceedings of the Workshop--Lisbon, 18 to 20 November 1993, Office for Official Publ. Eur. Communities, Luxembourg, pp. 69-81.

Liang, S. and A. H. Strahler, 1994, Retrieval of surface BRDF from multiangle remotely sensed data, Remote Sens. Environ., vol. 50, pp. 18-30.

Schaaf, C. B. and A. H. Strahler, 1994, Simulating the bidirectional and hemispherical reflectance of mountainous and forested scenes with a geometric-optical model, Proc. Seventh Conf. on Satellite Meteorology, June 6-10, Monterey, CA (American Meteorological Soc.), pp. 602-604.

Jupp, D. L. B., E. R. MacDonald, B. A. Harrison, X. Li, A. H. Strahler, and C. E. Woodcock, 1994, Prospects for mapping canopy structure using geometric-optical models, Proc. 7th Australasian Remote Sensing Conference, Melbourne, Australia, March 1-4, 1994, 9 pp.

Strahler, A. H., 1994, Vegetation Canopy Reflectance Modeling--Recent Developments and Remote Sensing Perspectives, Proc. Sixth Int. Symp. on Physical Measurements and Spectral Signatures, Val d'Isere, France, 17-21 January, 1994, pp. 593-600.

Liang, S. and A. H. Strahler, 1994, A stochastic radiative transfer model of a discontinuous vegetation canopy, Proc. 14th Int. Geosci. and Remote Sensing Symp., Pasadena, CA, Aug. 8-12, 1994, vol. 3, pp. 1626-1628.

Zhu, C.G., J. Wang and A. H. Strahler, 1994, A new registration algorithm for ASAS multi-angle images, Proc. 14th Int. Geosci. and Remote Sensing Symp., Pasadena, CA, Aug. 8-12, 1994, vol. 3, pp. 1658-1660.

Li, X., A. H. Strahler and C. E. Woodcock, 1994, A hybrid geometric optical-radiative transfer model for directional reflectance of discontinuous vegetation canopies, Proc. 14th Int. Geosci. and Remote Sensing Symp., Pasadena, CA, Aug. 8-12, 1994, vol. 3, pp. 1829-1831.

Schaaf, C. B., X. Li and A. H. Strahler, 1994, Validation of canopy bidirectional reflectance models with ASAS imagery of a spruce forest in Maine, Proc. 14th Int. Geosci. and Remote Sensing Symp., Pasadena, CA, Aug. 8-12, 1994, vol. 3, pp. 1832-1833.

Moody, A., S. Gopal, A. H. Strahler, J. Borak, and P. Fisher, 1994, A combination of temporal thresholding and neural network methods for classifying multiscale remotely-sensed image data, Proc. 14th Int. Geosci. and Remote Sensing Symp., Pasadena, CA, Aug. 8-12, 1994, vol. 4, pp. 1877-1879.

Strahler, A. H. and S. Liang, 1994, Comparisons of radiative transfer models of vegetation canopies with laboratory measurements, Proc. 14th Int. Geosci. and Remote Sensing Symp., Pasadena, CA, Aug. 8-12, 1994, vol. 4, pp. 2345-2347.

Liang, S., and A. H. Strahler, 1994, A four-stream solution for atmospheric radiance transfer over a non-Lambertian surface, Applied Optics, vol. 33, pp. 5745-5753.

Moody, A., and C. E. Woodcock, 1994, Scale-dependent errors in the estimation of land-cover proportions--Implications for global land-cover datasets, Photogram. Engr. and Remote Sens., vol. 60, pp. 585-594.

Lambin, E. F. and A. H. Strahler, 1994, Change-vector analysis: A tool to detect and categorize land-cover change processes using high temporal-resolution satellite data, Remote Sens. Environ., vol. 48, pp. 231-244.

Lambin, E. F. and A. H. Strahler, 1994, Indicators of land-cover change for change-vector analysis in multitemporal space at coarse spatial scales, Int. J. Remote Sens., vol. 15, pp. 2099-2119.

Abuelgasim, A. A. and A. H. Strahler, 1994, Modeling bidirectional radiance measurements collected by the Advanced Solid-State Array Spectroradiometer (ASAS) over Oregon Transect conifer forests, Remote Sens. of Environ., vol. 47, pp. 261-275.

Barnsley, M. J., A. H. Strahler, K. P. Morris, and J.-P. Muller, 1994, Sampling the surface bidirectional reflectance distribution function (BRDF): Evaluation of current and future satellite sensors, Remote Sensing Reviews, vol. 8, pp. 271-311.

Liang, S. and A. H. Strahler, 1993, Calculation of the angular radiance distribution for a coupled system of atmosphere and canopy media using an improved Gauss-Seidel algorithm, IEEE Trans. Geosci. and Remote Sens., vol. 31, pp. 491-502.

Schaaf, C. L. B., and A. H. Strahler, 1993, Modeling the bidirectional reflectance and spectral albedo of a conifer forest, Proceedings, 25th International Symposium on Remote Sensing and Global Environmental Change, April 2-8, 1993, Graz, Austria, vol. 2, pp. 594-601.

Schaaf, C. B. and A. H. Strahler, 1993, Solar zenith angle effects on forest canopy hemispherical reflectances calculated with a geometric-optical bidirectional reflectance model, IEEE Trans. Geosci. and Remote Sens., vol. 31, pp. 921-927.

Liang, S. and A. H. Strahler, 1993, An analytic BRDF model of canopy radiative transfer and its inversion, IEEE Trans. Geosci. and Remote Sens., vol. 31, pp. 1081-1092.

Li, X., and Strahler, A. H., 1992, Mutual shadowing and directional reflectance of a rough surface: A geometric-optical model, Proc. IGARRS '92, May 26-29, 1992, Houston, Texas, pp. 766-768.

Liang, S., and Strahler, A. H., 1992, An explicit canopy BRDF model and inversion, Proc. IGARRS '92, May 26-29, 1992, Houston, Texas, pp. 1487-1489.

Li, X. and A. H. Strahler, 1992, Geometric-optical bidirectional reflectance modeling of the discrete-crown vegetation canopy: Effect of crown shape and mutual shadowing, IEEE Trans. Geosci. and Remote Sens., vol. 30, pp. 276-292.

Jupp, D. L. B., B. A. Harrison, R. C. Gray, G. T. Byrne, E. R. McDonald, A. H. Strahler, C. E. Woodcock, and X. Li, 1992, Monitoring vegetation structure using geo-optical modelling and directional radiance, Proc. 13th Asian Conference on Remote Sensing, Spring, 1992.